arXiv:1407.3470v1 [math.RT] 13 Jul 2014

IRREDUCIBLE WEIGHT MODULES OVER WITT ALGEBRAS WITH INFINITE DIMENSIONAL WEIGHT SPACES

GENQIANG LIU AND KAIMING ZHAO

Abstract. Let $d > 1$ be an integer. In 1986, Shen defined a class of weight modules $F_b^\alpha(V)$ over the Witt algebra $\mathcal{W}_d$ for $\alpha \in \mathbb{C}^d$, $b \in \mathbb{C}$, and an irreducible module V over the special linear Lie algebra $\mathfrak{sl}_d$. In 1996, Eswara Rao determined the necessary and sufficient conditions for these modules to be irreducible when V is finite dimensional. In this note, we will determine the necessary and sufficient conditions for all these modules $F_b^\alpha(V)$ to be irreducible where V is not necessarily finite dimensional. Therefore we obtain a lot of irreducible $\mathcal{W}_d$-modules with infinite dimensional weight spaces.

Keywords: Witt algebra, weight module, irreducible module

2010 Math. Subj. Class.: 17B10, 17B20, 17B65, 17B66, 17B68

1. Introduction

We denote by $\mathbb{Z}$, $\mathbb{Z}_+$, $\mathbb{N}$ and $\mathbb{C}$ the sets of all integers, nonnegative integers, positive integers and complex numbers, respectively. Let $d > 1$ be an integer.

Representation theory for infinite-dimensional Lie algebras has been attracting extensive attentions of many mathematicians and physicists. These Lie algebras include the Witt algebras $\mathcal{W}_d$ which is the derivation algebra of the Laurent polynomial algebra $A_d = \mathbb{C}[x_1^{\pm 1}, x_2^{\pm 1}, \cdots, x_d^{\pm 1}]$. The algebra $\mathcal{W}_d$ is a natural higher rank generalization of the Virasoro algebra, which has many applications to different branches of mathematics and physics (see [M, L1, L2, L3, L4, L5]) and at the same time a much more complicated representation theory.

The weight representation theory of Witt algebras was recently studied by many experts; see [B, E1, E2, BMZ, GLZ, L3, L4, L5, MZ2, Z]. In 1986, Shen defined a class of modules $F_b^\alpha(V)$ over the Witt algebra $\mathcal{W}_d$ for $\alpha \in \mathbb{C}^d$, $b \in \mathbb{C}$, and an irreducible module V over the special linear Lie algebra $\mathfrak{sl}_d$, see [Sh], which were also given by Larsson in 1992, see [L3]. In 1996, Eswara Rao determined the necessary and sufficient conditions for these modules to be irreducible when V is finite dimensional, see [E1]. A simplified proof was given in [GZ]. It is natural to study the irreducibility of these modules $F_b^\alpha(V)$ when V is infinite dimensional. This is what the present paper will do. In this

manner we obtain a lot of irreducible modules over the Witt algebra $\mathcal{W}_d$ with infinite dimensional weight spaces.

We have to mention the remarkable work by Billig and Futorny [BF] in which they proved very recently that irreducible modules for $\mathcal{W}_d$ with finite-dimensional weight spaces fall in two classes: (1) modules of the highest weight type and (2) irreducible modules from $F_b^\alpha(V)$. So the next task in representation theory of Witt algebras is to study irreducible modules over the Witt algebra $\mathcal{W}_d$ with infinite dimensional weight spaces, and irreducible non-weight modules. This is the second reason of this paper.

For irreducible weight modules (not necessary with finite dimensional weight spaces) over $\mathcal{W}_d$, all weight sets were explicitly determined in [MZ2], while some non-weight irreducible $\mathcal{W}_d$-modules were constructed in [TZ].

In this paper we actually prove that, for any infinite dimensional irreducible $\mathfrak{sl}_d$-module V, any $\alpha \in \mathbb{C}^d$ and $b \in \mathbb{C}$, the $\mathcal{W}_d$-module $F_b^\alpha(V)$ is always irreducible, see Theorem 2.4. We also determine the necessary and sufficient conditions for two $\mathcal{W}_d$-module $F_b^\alpha(V)$ to be isomorphic, see Theorem 2.6. The main trick we employed in our proof is a characterization of finite dimensional irreducible modules over $\mathfrak{sl}_d$, see Lemma 2.3, which makes the proof very short, and very different from that for finite dimensional irreducible $\mathfrak{sl}_d$-module V, see [E1, GZ].

Note that $F_b^\alpha(V)$ are polynomial modules in the sense of [BB, BZ].

2. $\mathcal{W}_d$-modules from $\mathfrak{sl}_d$-modules

For a positive integer $d > 1$, $\mathbb{Z}^d$ denotes the direct sum of d copies of $\mathbb{Z}$. For any $a = (a_1, \cdots, a_d)^T \in \mathbb{Z}_+^d$ and $n = (n_1, \cdots, n_d)^T \in \mathbb{C}^d$, we denote $n^a = n_1^{a_1} n_2^{a_2} \cdots n_d^{a_d}$, where T means taking transpose of the matrix. Let $\mathfrak{gl}_d$ be the Lie algebra of all $d \times d$ complex matrices, $\mathfrak{sl}_d$ the subalgebra of $\mathfrak{gl}_d$ consisting of all traceless matrices. For $1 \leqslant i, j \leqslant d$ we use E_{ij} to denote the matrix units.

2.1. **Witt algebras $\mathcal{W}_d$.** We denote by $\mathcal{W}_d$ the derivation Lie algebra of the Laurent polynomial algebra $A_d = \mathbb{C}[x_1^{\pm 1}, x_2^{\pm 1}, ..., x_d^{\pm 1}]$. For $i \in \{1, 2, \ldots, d\}$, set $\partial_i = x_i \frac{\partial}{\partial x_i}$; and for any $a = (a_1, a_2, \cdots, a_d)^T \in \mathbb{Z}^d$ (considered as column vectors), set $x^a = x_1^{a_1} x_2^{a_2} \cdots x_d^{a_d}$.

We fix the vector space $\mathbb{C}^d$ of $d \times 1$ matrices. Denote the standard basis by $\{e_1, e_2, ..., e_d\}$. Let $(\cdot\,|\,\cdot)$ be the standard symmetric bilinear form such that $(u|v) = u^T v \in \mathbb{C}$. For $u \in \mathbb{C}^d$ and $r \in \mathbb{Z}^d$, we denote $D(u, r) = x^r \sum_{i=1}^d u_i \partial_i$. Then we have

$$[D(u, r), D(v, s)] = D(w, r + s),\ u, v \in \mathbb{C}^d, r, s \in \mathbb{Z}^d,$$

where $w = (u|s)v - (v|r)u$. Note that for any $u, v, z, y \in \mathbb{C}^d$, both uv^T and xy^T are $d \times d$ matrices, and

$$(uv^T)(zy^T) = (v|z)uy^T.$$

We know that $\mathfrak{H} = \text{span}\{\partial_1, \partial_2, ..., \partial_d\}$ is the Cartan sunalgebra of $\mathcal{W}_d$. A weight $\mathcal{W}_d$-module is a $\mathcal{W}_d$-module V so that $V = \oplus_{\lambda \in \mathfrak{H}^*} V_\lambda$ where

$$V_\lambda = \{v \in V : hv = \lambda(h)v \ \forall \ h \in \mathfrak{H}\}.$$

2.2. **Defining $\mathcal{W}_d$-modules.** Let us first recall Shen's Witt modules from [Sh]. For any $\alpha \in \mathbb{C}^d, b \in \mathbb{C}$ and $\mathfrak{gl}_d$-module V on which the identity matrix acts as the scalar b, let $F_b^\alpha(V) = V \otimes A_d$. For simplicity we write $v(n) = v \otimes x^n$ for any $v \in V, n \in \mathbb{Z}^d$. Then $F_b^\alpha(V)$ becomes a $\mathcal{W}_d$-module if we define the following actions

$$D(u,r)v(n) = \Big((u \mid n+\alpha)v + (ru^T)v\Big)(n+r), \tag{2.1}$$

where $u \in \mathbb{C}^d$, $v \in V$, $n, r \in \mathbb{Z}^d$. It is easy to see that the module $F_b^\alpha(V)$ obtained from any V is always a weight module over $\mathcal{W}_d$.

When V is finite dimensional, the following Theorem was given by Eswara Rao, see [E2] or [GZ].

t **Theorem 2.1.** *Let $\alpha \in \mathbb{C}^d, b \in \mathbb{C}$, and V be a finite dimensional irreducible module over $\mathfrak{gl}_d$ on which the identity matrix acts as the scalar b. Then $F_b^\alpha(V)$ is a reducible $\mathcal{W}_d$-module if and only if the highest weight of V is the fundament weight ω_k of $\mathfrak{gl}_d$ and $b = k$, where $k \in \mathbb{Z}$ with $1 \leqslant k \leqslant d-1$.*

2.3. **Description of $F_b^\alpha(V)$.** The following useful lemma from linear algebra is obvious.

l **Lemma 2.2.** *For $n = (n_1, n_2, ..., n_d), u = (u_1, u_2, ..., u_d) \in \mathbb{C}^d$, let*

$$g(n,u) = \sum_{a,b \in \mathbb{Z}_+^d} c_{a,b} n^a u^b$$

be a polynomial in $2d$ variables n_i, u_j with coefficients $c_{a,b} \in \text{End}(V)$. Let V^* be a subspace of V. For $v \in V$, if $g(n,u)v \in V^*$ for all $n \in \mathbb{Z}^d, u \in \mathbb{C}^d$, then $c_{a,b}v \in V^*$ for any $a, b \in \mathbb{Z}_+^d$.

Now we need the following characterization on finite dimensional irreducible $\mathfrak{sl}_d$-modules.

l2 **Lemma 2.3.** *Let V be an irreducible $\mathfrak{sl}_d$-module(not necessarily weight module).*

1. *For any $i, j : 1 \leqslant i \neq j \leqslant d$, E_{ij} acts injectively or locally nilpotently on V.*
2. *The module V is finite dimensional iff E_{ij} acts locally nilpotently on V for any $i, j : 1 \leqslant i \neq j \leqslant d$.*

Proof. (1). If E_{ij} is not injective on V, then $E_{ij}v = 0$ for some nonzero vector v in V. It is known that $\text{ad}E_{ij}$ is locally nilpotent on the universal enveloping algebra $U(\mathfrak{sl}_d)$ of $\mathfrak{sl}_d$. This means that, for any

$X \in U(\mathfrak{sl}_d)$ we have $(\mathrm{ad}E_{ij})^k X = 0$ for sufficiently large k. Then from the identity

$$E_{ij}^k Xv = ((\mathrm{ad}E_{ij})^k X)v = 0,$$

for sufficiently large k. Since V is irreducible, $V = U(\mathfrak{sl}_d)v$. Thus E_{ij} acts locally nilpotently on V.

(2). If V is finite dimensional, clearly E_{ij} acts locally nilpotently on V for any $i, j : 1 \leqslant i \neq j \leqslant d$.

Now suppose that E_{ij} acts locally nilpotently on V for any $i, j : 1 \leqslant i \neq j \leqslant d$. By the condition that E_{ij} acts locally nilpotently on V for any $i, j \in \mathbb{Z}$ with $1 \leqslant i < j \leqslant d$ and using Theorem 1 in [MZ1], we see that V is a highest weight module. By the condition that E_{ij} acts locally nilpotently on V for any $i, j \in \mathbb{Z}$ with $1 \leqslant j < i \leqslant d$ and using the same Theorem, we can see that V is a lowest weight module. Thus V is finite dimensional. □

Now we are ready to prove our main result in this paper.

Theorem 2.4. *Let V be an infinite dimensional irreducible module over $\mathfrak{gl}_d$ on which the identity matrix acts as a scalar $b \in \mathbb{C}$. Then $F_b^\alpha(V)$ is an irreducible module over $\mathcal{W}_d$ for any $\alpha \in \mathbb{C}^d$.*

Proof. From Lemma 2.3 we know that the action E_{st} on V is injective for some $s, t : 1 \leq s \neq t \leq d$.

Suppose W is a nonzero $\mathcal{W}_d$-submodule of $F_b^\alpha(V)$. Next we will show that $W = F_b^\alpha(V)$. Since $F_b^\alpha(V)$ is a weight module over $\mathcal{W}_d$, then $W = \oplus_{n\in\mathbb{Z}^d} M_n \otimes t^n$, where W_n are subspaces of V. Let $W^* = \bigcap_{n\in\mathbb{Z}^d} W_n$. We will show that W^* is a nonzero $\mathfrak{gl}_d$-submodule of V.

Claim 1. W^* is nonzero.

Since W is nonzero, without loss of generality, we assume $W_0 \neq 0$. Choose a nonzero vector $v \in W_0$, fix some $m \in \mathbb{Z}^d$. For any $n \in \mathbb{Z}^d, u \in \mathbb{C}^d$, we have

$$\begin{aligned}
&D(u, m-n)D(u, n)v(0)\\
=&\Big((u \mid \alpha+n) + (m-n)u^T\Big)\Big((u \mid \alpha) + nu^T\Big)v(m),\\
=&\Big((u \mid \alpha+n) + \sum_{i,j}(m_i - n_i)u_j E_{ij}\Big)\Big((u \mid \alpha) + \sum_{k,l} n_k u_l E_{kl}\Big)v(m),\\
=&\Big(\sum_{i,j}\sum_{k,l}(m_i - n_i)u_j n_k u_l E_{ij}E_{kl} + (u \mid \alpha)\sum_{i,j}(m_i - n_i)u_j E_{ij}\\
&+ (u \mid \alpha+n)\sum_{k,l} n_k u_l E_{kl} + (u \mid \alpha)(u \mid n+\alpha)\Big)v(m)\\
=&\Big(\sum_{i,j}\sum_{k,l}(m_i - n_i)u_j n_k u_l E_{ij}E_{kl} + \sum_{i,j}((u \mid \alpha)m_i + (u \mid n)n_i)u_j E_{ij}\\
&+ (u \mid \alpha)(u \mid n+\alpha)\Big)v(m).
\end{aligned}$$

Set

$$\begin{aligned} g(n,u) =& \sum_{i,j}\sum_{k,l}(m_i - n_i)u_j n_k u_l E_{ij}E_{kl} \\ &+ \sum_{i,j}((u \mid \alpha)m_i + (u \mid n)n_i)u_j E_{ij} \\ &+ (u \mid \alpha)(u \mid n+\alpha) \end{aligned}$$

which is a polynomial in $n_i, u_j, 1 \leqslant i, j \leqslant d$ with coefficients in $\mathrm{End}(V)$. Then $g(n,u)v \in W_m$ for any $n \in \mathbb{Z}^d, u \in \mathbb{C}^d$.

Since $s \neq t$, the coefficient of $n_s^2 u_t^2$ in $g(n,u)$ is $-E_{st}E_{st}$. By Lemma 2.2, we can deduce that $E_{st}E_{st}v \in W_m$ for any $m \in \mathbb{Z}^d$. Thus $E_{st}E_{st}v \in W^*$. By the fact that the action E_{st} on V is injective, we see that $W^* \neq 0$. Claim 1 follows.

Claim 2. W^* is a $\mathfrak{gl}_d$-submodule of V.

By the definition of W^*, we see that $v(m-n) \in W_{m-n} \otimes x^{n-m}$ for any $n, m \in \mathbb{Z}^d$, $v \in W^*$. For any $n \in \mathbb{Z}_+^d, u \in \mathbb{C}^n$ we compute that

$$\begin{aligned} & D(u,n)v(m-n) \\ =& \Big((u \mid \alpha + m - n) + nu^T)v(m), \\ =& ((u \mid \alpha + m) - \sum_i u_i n_i + \sum_{i,j} n_i u_j E_{ij})v(m) \in W_m \otimes x^m. \end{aligned}$$

Considering the coefficient of $n_i u_j$ in the right hand of the last equality and using Lemma 2.2 again, we know that $E_{ij}v \in W_m$ for all $m \in \mathbb{Z}^d$ and $i, j : 1 \leqslant i, j \leqslant d$. This implies that $E_{ij}v \in W^*$ for all $v \in W^*$ and $i, j : 1 \leqslant i, j \leqslant d$. Therefore W^* is a $\mathfrak{gl}_d$-submodule of V.

By Claim 1 and Claim 2, we deduce that $W^* = V$. Therefore $W = F_b^\alpha(V)$, hence $F_b^\alpha(V)$ is an irreducible module over $\mathcal{W}_d$. □

Combining Theorem 2.4 and Theorem 2.1, we obtain the following:

Theorem 2.5. *Let $\alpha \in \mathbb{C}^d, b \in \mathbb{C}$, and V be an irreducible module over $\mathfrak{gl}_d$ on which the identity matrix acts as the scalar b. Then $F_b^\alpha(V)$ is a reducible module over $\mathcal{W}_d$ if and only if V is isomorphic to the finite dimensional module whose highest weight is the fundament weight ω_k and $b = k$, where $k \in \mathbb{Z}$ with $1 \leqslant k \leqslant d-1$.*

Next we will give a concrete example of $F_b^\alpha(V)$ for which V is infinite dimensional.

Example. In [N], Nilsson constructed a class of irreducible non-weight modules over $\mathfrak{sl}_d$. Explicitly, for $1 \leqslant k \leqslant d-1$, let $h_k = E_{k,k} - \frac{1}{d}\sum_{i=1}^{d} E_{ii}$. Then for each $\beta \in \mathbb{C}$, the polynomial algebra $\mathbb{C}[h_1, \ldots, h_{d-1}]$ is an

irreducible $\mathfrak{sl}_d$-module under the action

$$\begin{aligned}
h_i \cdot f(h_1, \dots, h_{d-1}) &= h_i f(h_1, \dots, h_{d-1}),\\
E_{i,d} \cdot f(h_1, \dots, h_{d-1}) &= (\beta + \sum_{k=1}^{d-1} h_k)(h_i - \beta - 1) f(h_1, \dots, h_i - 1, \dots, h_{d-1}),\\
E_{d,j} \cdot f(h_1, \dots, h_{d-1}) &= -f(h_1, \dots, h_j + 1, \dots, h_{d-1}),\\
E_{i,j} \cdot f(h_1, \dots, h_{d-1}) &= (h_i - \beta - 1) f(h_1, \dots, h_i - 1, \dots, h_j + 1, \dots, h_{d-1}),
\end{aligned}$$

where $1 \leqslant i, j \leqslant d-1$. By Theorem 2.4, $F_b^\alpha(\mathbb{C}[h_1, \dots, h_{d-1}])$ is an irreducible weight $\mathcal{W}_d$-module with infinite dimensional weight spaces.

Finally, we can consider the isomorphism between two modules described above.

Theorem 2.6. *Let $\alpha, \beta \in \mathbb{C}^d$, $b, b' \in \mathbb{C}$, and V, W be two irreducible $\mathfrak{sl}_d$-modules. Then the $\mathcal{W}_d$-modules $F_b^\alpha(V)$ and $F_{b'}^\beta(V')$ are isomorphic if and only if $b = b', \alpha - \beta \in \mathbb{Z}^d$ and $V \cong V'$ as $\mathfrak{sl}_d$-modules.*

Proof. The sufficiency of the condition is clear. Now suppose that $\rho : F_b^\alpha(V) \to F_{b'}^\beta(V')$ a $\mathcal{W}_d$-module isomorphism. Since isomorphic modules have the same weight set, we have $\alpha + \mathbb{Z}^d = \beta + \mathbb{Z}^d$, i.e., $\alpha - \beta \in \mathbb{Z}^d$. Without loss of generality, we assume $\alpha = \beta$. Then ρ induces a linear isomorphism σ from V to V' such that

$$\rho(v(0)) = \sigma(v)(0), \quad \forall\ v \in\ V.$$

From $\rho(D(u,n)v(0)) = D(u,n)\rho(v(0))$, we see that

$$\sum_{i,j} n_i u_j \sigma(E_{ij} v) = \sum_{i,j} n_i u_j E_{ij} \sigma(v) \tag{2.2}$$

for all $n \in \mathbb{Z}^d, u \in \mathbb{C}^d$. Comparing the coefficient of $n_i u_j$ on both sides of the formula (2.2), we know that $\sigma(E_{ij} v) = E_{ij}\sigma(v)$ for any $v \in V'$ and $1 \leqslant i, j \leqslant d$. Therefore $V \cong V'$ as $\mathfrak{gl}_d$-modules. □

Acknowledgments

G.L. is partially supported by NSF of China (Grant 11301143) and the school fund of Henan University(2012YBZR031, 0000A40382); K.Z. is partially supported by NSF of China (Grant 11271109) and NSERC.

References

[B] Y. Billig, Jet modules, Canad. J. Math., 59 (2007), no.4, 712-729.

[BB] S. Berman and Y. Billig, Irreducible representations for toroidal Lie algebra, *J. Algebra*, 221(1999), 188-231

[BF] Y. Billig and V. Futorny, Classification of simple W_n-modules with finite-dimensional weight spaces, preprint, arXiv:1304.5458v1.

[BMZ] Y. Billig, A. Molev, R. Zhang, Differential equations in vertex algebras and simple modules for the Lie algebra of vector fields on a torus, Adv. Math., 218(2008), no.6, 1972-2004.

[BZ] Y. Billig and K. Zhao, Weight modules over exp-polynomial Lie algebras, J. Pure Appl. Algebra, 191(2004), 23-42

[DMP] I. Dimitrov, O. Mathieu, I. Penkov, On the structure of weight modules. Trans. Amer. Math. Soc. 352 (2000), no. 6, 2857-2869.

[E1] S. Eswara Rao, Irreducible representations of the Lie-algebra of the diffeomorphisms of a d-dimensional torus, J. Algebra, **182** (1996), no. 2, 401–421.

[E2] S. Eswara Rao, Partial classification of modules for Lie algebra of diffeomorphisms of d-dimensional torus, J. Math. Phys., 45 (8), (2004) 3322-3333.

[GLZ] X. Guo, G. Liu and K. Zhao, Irreducible Harish-Chandra modules over extended Witt algebras, Ark. Mat., 52 (2014), 99-112.

[GZ] X.Guo and K.Zhao, Irreducible weight modules over Witt algebras, Proc. Amer. Math. Soc., 139(2011), 2367-2373.

[L1] T. A. Larsson, Multi dimensional Virasoro algebra, Phys. Lett., B 231, 94-96(1989).

[L2] T. A. Larsson, Central and non-central extensions of multi-graded Lie algebras, J. Phys., A 25, 1177-1184(1992).

[L3] T. A. Larsson, Conformal fields: A class of representations of Vect (N), Int. J. Mod. Phys., A 7, 6493-6508(1992).

[L4] T. A. Larsson, Lowest energy representations of non-centrally extended diffeomorphism algebras, Commun. Math. Phys., 201, 461-470(1999).

[L5] T. A. Larsson, Extended diffeomorphism algebras and trajectories in jet space, Commun. Math. Phys., 214, 469-491(2000).

[M] O. Mathieu; Classification of irreducible weight modules. Ann. Inst. Fourier (Grenoble) 50 (2000), no. 2, 537-592.

[MZ1] V. Marzuchuk, K. Zhao, Characterization of simple highest weight modules. Can. Math. Bull. 56(3), 606-614 (2013)

[MZ2] V. Marzuchuk and K. Zhao, Supports of weight modules over Witt algebras, Proc. Roy. Soc. Edinburgh Sect., A 141(2011), no. 1, 155-170.

[N] J. Nilsson, Simple $\mathfrak{sl}_{n+1}$-module structures on $U(h)$, arXiv:1312.5499.

[Sh] G. Shen, Graded modules of graded Lie algebras of Cartan type. I. Mixed products of modules, Sci. Sinica Ser., A **29** (1986), no. 6, 570-581.

[TZ] H. Tan, K. Zhao, $\mathcal{W}_n^+$ and $\mathcal{W}_n$-module structures on $U(h)$, arXiv:1401.1120.

[Z] K. Zhao, Weight modules over generalized Witt algebras with 1-dimensional weight spaces, Forum Math., Vol.16, No.5, 725-748.

G.L.: College of Mathematics and Information Science, Henan University, Kaifeng 475004, China. Email: liugenqiang@amss.ac.cn

K.Z.: Department of Mathematics, Wilfrid Laurier University, Waterloo, ON, Canada N2L 3C5, and College of Mathematics and Information Science, Hebei Normal (Teachers) University, Shijiazhuang, Hebei, 050016 P. R. China. Email: kzhao@wlu.ca